

Solutions of $F(u, u_t, u_{x_1}, \dots, u_{x_n}) = 0$

S. Y. Lou^{1,2*}

¹ Shanghai Key Laboratory of Trustworthy Computing, East China Normal University, Shanghai 200062, China

² Faculty of Science, Ningbo University, Ningbo, 315211, China

Abstract: All solutions of $F(u, u_t, u_{x_1}, \dots, u_{x_n}) = 0$ are determined by its symmetries $\sigma = G(\tau_1, \dots, \tau_n)$ with $\tau_i \equiv u_{x_i}/u_t$ and a seed solution set. Especially, the Lax pair, recursion operator, group invariant operator, infinitely many high order symmetries and higher order group invariant functions are also explicitly given for arbitrary (1+1)-dimensional first order partial differential equations.

It is known that symmetry theory is powerful to find exact solutions of any physical systems [1]. The singularity analysis (Painlevé analysis) shows us that if a partial differential equation (PDE) system is Painlevé integrable, then enough arbitrary functions should be included in a formal series solution with respect to an arbitrary singularity manifold [2]. Thus, if one tries to find general solutions of a PDE system by means of symmetry theory, then one has to find enough symmetries with enough arbitrary functions. Inversely, if you can find enough symmetries with enough arbitrary functions, then, you may successful find all the solutions of the related PDE system. To realized this idea, the $(n+1)$ -dimensional first order PDE

$$F(u, u_t, u_{x_1}, \dots, u_{x_n}) = 0, \quad (1)$$

is an excellent example.

According to this idea, to find the general solution of (2) is equivalent to find a symmetry including one (one is enough for a first order PDE) n -dimensional arbitrary function, say,

$$\sigma = \sigma(t, x_1, \dots, x_n, u, G(\tau_1, \dots, \tau_n)), \quad (2)$$

where G is an arbitrary function of n independent variables $\tau_i = \tau_i(t, x_1, \dots, x_n)$.

A symmetry of (2) σ is defined as a solution of its linearized equation

$$F'\sigma \equiv \left(\frac{\partial F}{\partial u} + \frac{\partial F}{\partial u_t} \partial_t + \sum_{i=1}^n \frac{\partial F}{\partial u_{x_i}} \partial_{x_i} \right) \sigma = 0 \quad (3)$$

which means (2) is invariant under the transformation

$$u \rightarrow u + \epsilon \sigma$$

*Email: lousenyue@nbu.edu.cn

with infinitesimal ϵ .

Fortunately, by substituting

$$\sigma = G(\tau_1, \dots, \tau_n)u_t, \quad \tau_i = \frac{u_{x_i}}{u_t}, \quad (4)$$

into (3), one can readily prove that σ given by (4) with G being an arbitrary function is really a symmetry of (2).

Now, the remained problem is how to use the symmetry to find the general solution of (2). Because of the entrance of u_{x_i} , $i = 1, \dots, n$ and u_t in the symmetry (4), we have to study the related transformations of u_{x_i} and u_t . Differentiating (4) with respect to x_i and t , it is straightforward to find

$$\sigma^{u_i} = \left(G - \sum_{j=1}^n \tau_j G_{\tau_j} \right) u_{it} + \sum_{j=1}^n G_{\tau_j} u_{ix_j}, \quad u_i \equiv u_{x_i}, \quad (5a)$$

$$\sigma^{u_t} = \left(G - \sum_{j=1}^n \tau_j G_{\tau_j} \right) u_{tt} + \sum_{j=1}^n G_{\tau_j} u_{tx_j}. \quad (5b)$$

The above result (5) leads us to rewrite the symmetry (4) as

$$\sigma = \left(G - \sum_{j=1}^n \tau_j G_{\tau_j} \right) u_t + \sum_{j=1}^n G_{\tau_j} u_{x_j} \equiv T u_t + \sum_{j=1}^n X_j u_{x_j}. \quad (6)$$

From (6) and (5), we know that the symmetry (4) can be equivalently considered as the pure space-time transformations with the vector

$$V = T \partial_t + \sum_{j=1}^n X_j \partial_{x_j}. \quad (7)$$

It is fortunate that the first order prolongations of the vector related to u_{x_i} and u_t are all vanished. Thus u , iu_{x_i} and u_t and then τ_i are all group invariants. This fact leads to the following non-degeneration solution theorem.

Theorem 1. *Non-degeneration solution theorem.* If $u_0(t, x_i, i = 1, 2, \dots, n)$ is a non-degenerate solution of (2), so u is with

$$u = u_0(t(\epsilon), x_i(\epsilon), i = 1, 2, \dots, n) \quad (8)$$

while $t(\epsilon)$, $x_i(\epsilon)$, $i = 1, 2, \dots, n$ are determined by

$$t(\epsilon) = t - \epsilon \left[G(\eta) - \sum_{j=1}^n \eta_j G_{\eta_j} \right], \quad (9a)$$

$$x_i(\epsilon) = x_i - \epsilon G_{\eta_i}, \quad \eta_j \equiv \frac{u_{0x_j}}{u_{0t}} \Big|_{t \rightarrow t(\epsilon), x_i \rightarrow x_i(\epsilon), i=1, \dots, n}. \quad (9b)$$

The definition of the non-degeneration solution is defined by η_i , $i = 1, \dots, n$ (and then τ_i) in the solution (12) are independent.

Because the model (2) is $(n+1)$ -dimensional, the arbitrary functions included in its general solutions should be n -dimensional. If η_j , $j = 1, \dots, n$ are degenerate, the Theorem 1 can still be used to find special

solutions of the model by just taking away the terms with ∂_{x_j} if $\eta_j = \text{const.}$ (special trivial degenerated cases) in (9).

However, to find all the degenerate solutions of the model equation, we have to find all the possible non-degenerate solutions in lower dimensions. In degenerate cases, without loss of generality, we can express $\tau_n = u_{x_n}/u_t$ by

$$u_{x_n} = H(\tau_1, \dots, \tau_{n-1})u_t. \quad (10)$$

Substituting (10) into (2) yields an $(m+1)$ -dimensional system with $m = n-1$. Using the Theorem 1 to $m = n-1$ case, we can find all possible non-degenerate solutions of the model in n dimensional space-time. Repeated the step, we can find all the possible degenerate solutions of the model in all the lower dimensions.

As the last step to find all the possible degenerate solutions of the model (2), we should investigate in detail for the $n=1$ case of (2) with $x_1 \equiv x$. In (1+1)-dimensional case, I write (2) in its equivalent form

$$u_t = F(u, u_x). \quad (11)$$

For simplicity, I just list all the results I have obtained for the model (11).

Theorem 2. If $u_0(x, t)$ is a non-traveling wave solution of (11), so $u = u(x, t)$ is with

$$u = u(x, t) = u_0(t(\epsilon), x(\epsilon)) \quad (12)$$

while $t(\epsilon)$ and $x(\epsilon)$ are determined by

$$t(\epsilon) = t - \epsilon [G(\eta) - \eta G_\eta], \quad (13a)$$

$$x(\epsilon) = x - \epsilon G_\eta, \quad \eta \equiv \frac{u_{0x}}{u_{0t}} \Big|_{t \rightarrow t(\epsilon), x \rightarrow x(\epsilon)}. \quad (13b)$$

Theorem 2 is only the special case of the Theorem 1 because the traveling wave solution is the only degenerate solution of (11). Obviously, the degenerate solution, the traveling wave solution, of (1) can be obtained by once integration with respect to the traveling variable $\xi \equiv x - ct$.

The (1+1)-dimensional model possesses some recursion operators, say,

$$\Phi = \frac{u_x K_1}{u_{xx} + K_2} \partial_x u_x^{-1} \quad (14)$$

where $K_i = K_i(u, u_x) \equiv K_i(y_0, y_1)$, $i = 1, 2$ are defined as

$$y_1 F_{y_0} K_{2y_1} + (F - y_1 F_{y_1}) K_{2y_0} + F_{y_1 y_1} K_2^2 - (2y_1 F_{y_0 y_1} + F_{y_0}) K_2 + y_1^2 F_{y_0 y_0} = 0, \quad (15a)$$

$$y_1 F_{y_0} K_{1y_1} + (F - y_1 F_{y_1}) K_{1y_0} + F_{y_1 y_1} K_2 K_1 - (y_1 F_{y_0 y_1} + F_{y_0}) K_2 = 0. \quad (15b)$$

The eigenvalue problem of the recursion operator Φ and the symmetry definition equation (3) constitutes a weak Lax pair of the model (11).

If we define the the group invariant function (GIF), I , as a solution of

$$I_t = F_{u_x} I_x, \quad (16)$$

and σ_0 is a known symmetry, then

$$\sigma = I\sigma_0 \quad (17)$$

is also a symmetry of (11). The similar conclusion is hold for the general case (2).

For any GIF I , its arbitrary function $J(I)$ is also a GIF. Similar to the symmetry case, there are also some types of recursion operators (group invariant operators) which transform any GIF I_0 to a new GIF, say,

$$I = \phi I_0 = \frac{K_1}{u_{xx} + K_2} \partial_x I_0 \quad (18)$$

with the same K_i as defined in (15).

Using the group invariant operator(s), we have the following symmetry theorem for the (1+1)-dimensional system (11).

Theorem 3. *Symmetry theorem.* The (1+1)-dimensional system (11) possesses infinitely many higher order symmetries

$$\sigma = G \left(\phi^i \frac{u_x}{u_t}, \phi^j V, \phi^k W, i = 1, 2, \dots, n_1, j = 1, 2, \dots, n_2, k = 1, 2, \dots, n_3 \right) u_t, \quad (19)$$

where n_1, n_2 and n_3 are arbitrary integers, ϕ is a group invariant operator, say, that defined by (18), V and W are defined by

$$V = x + \int^u \frac{F_y(b, y(u, u_x, b)) db}{y(u, u_x, b)F_y(b, y(u, u_x, b)) - F(b, y(u, u_x, b))}, \quad (20a)$$

$$W = t + \int^u \frac{db}{F(b, y(u, u_x, b)) - y(u, u_x, b)F_y(b, y(u, u_x, b))}, \quad (20b)$$

with $y \equiv y(u, u_x, b)$ being determined by

$$F(u, u_x)y - u_x F(b, y) = 0. \quad (21)$$

Because the space-time variables x and t are in the equivalent position, all the results related the model (11) are still valid by exchanging the variables x and t .

Finally, we apply the above general theory to the following special example,

$$u_t = uu_x^2. \quad (22)$$

For the special model (22), one of the recursion operators has the form

$$\Phi_1 = u_x \phi_1 u_x^{-1} \quad (23)$$

with ϕ_1 being a group invariant operator

$$\phi_1 = u^{-3} u_{xx}^{-1} \partial_x. \quad (24)$$

Some special higher order symmetries can be expressed by

$$\sigma = G \left(\phi_1^i \frac{u_x}{u_t}, \phi_1^j (x - uu_x^{-1}), \phi_1^k (2t + u_x^{-2}), i = 1, 2, \dots, n_1, j = 1, 2, \dots, n_2, k = 1, 2, \dots, n_3 \right) u_t, \quad (25)$$

with G being an arbitrary function of the indicated variables.

The model (22) possesses a special non-traveling wave solution

$$u_0 = \frac{x}{\sqrt{-2t}} \quad (26)$$

which can be directly verified or derived from a high order (second order) symmetry constraint.

By using the Theorem 2 and the special non-degenerate solution (26), the general solution of the toy model (22) has the form

$$u = \frac{\xi}{\sqrt{-2\tau}}, \quad (27a)$$

$$u = \sqrt{2cx + 2c^2t + c_0}, \quad (27b)$$

where c and c_0 are arbitrary constants and $\xi = \xi(x, t)$ and $\tau = \tau(x, t)$ are determined by

$$\xi = x - \epsilon G_\eta, \quad \eta \equiv -2\tau\xi^{-1}, \quad (28a)$$

$$\tau = t - \epsilon(G + 2\tau\xi^{-1}G_\eta), \quad (28b)$$

with $G \equiv G(\eta)$ being an arbitrary function of η .

The special solution expressed in (28b) is the only degenerate solution, traveling wave solution of the model (22).

In summary, arbitrary first order PDEs are C-integrable. All the solutions of them can be obtained from the same solution theorem 1. The only model dependent information is included in their non-degenerate seed solutions. In (1+1)-dimensional case, many other interesting properties such as the Lax pair, recursion operator, group invariant operator, higher order symmetries and group invariant functions are also explicitly given. The similar results for higher dimensional first order PDEs can also be obtained in a similar way.

The author is grateful to thank Professors CW Cao, RX Yao, YQ Li, Y Chen, QP Liu, XB Hu, EG Fan and DJ Zhang for their helpful suggestions and fruitful discussions. The work was sponsored by the National Natural Science Foundations of China (Nos. 11175092, 11275123, 11205092 and 10905038), Shanghai Knowledge Service Platform for Trustworthy Internet of Things (No. ZF1213) and K. C. Wong Magna Fund in Ningbo University.

References

- [1] Olver P J, Application of Lie Groups to Differential Equation, Graduate Texts in Mathematics, 2nd ed. (Springer-Verlag, NewYork,1993).
- [2] Weiss J, Tabor M and Carnevale G 1983 J. Math. Phys. **24** 522
 Ramani A, Grammaticos B and Bountis T 1989 Phys. Rep. **180** 159
 Conte R 1989 Phys. Lett. A **140** 383
 Lou S Y 1998 Z. Naturforsch A **53** 51